



Anti-c-Myc tag antibody

Catalog: RGM2030


Product Information

Description:	Rabbit polyclonal antibody
Synonyms:	c-Myc
Immunogen:	Synthetic peptide of fragment c-Myc.
Form:	Liquid, 20 μ l, 1mg/ml
Purification:	Protein A purified
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 $^{\circ}$ C as supplied. 1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^{\circ}$ C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Confirmed Reactivity:	Anti-c-Myc recognizes c-Myc-tagged fusion proteins.

Application Example



Lane 1: 2.5 ng c-Myc-tagged fusion protein.

Lane 2: 5 ng c-Myc-tagged fusion protein.

Lane 3: 10 ng c-Myc-tagged fusion protein.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4 $^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 $^{\circ}$ C.

Secondary antibody: 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# HGR1020)

Detection: using chemiluminescence substrate and image were captured with CCD camera.